



Collection of Emission from an Oscillating Dipole Inside a Sphere: Analytic Integration Over a Circular Aperture

by J. David Pendleton and Steven C. Hill

ARL-TR-1399

December 1997

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof.

Destroy this report when it is no longer needed. Do not return it to the originator.

Army Research Laboratory

Adelphi, MD 20783-1197

ARL-TR-1399

December 1997

Collection of Emission from an Oscillating Dipole Inside a Sphere: Analytic Integration Over a Circular Aperture

J. David Pendleton and Steven C. Hill

Information Science and Technology Directorate

Abstract

We describe a method for analytically integrating, over a circular aperture, the emission from an oscillating dipole inside a dielectric sphere. The model is useful for investigating fluorescence, Raman, or other emission from molecules inside spherical particles or droplets. The analysis is performed for two cases: (1) the dipole emits from a fixed orientation, and (2) the dipole emits from all orientations. In both cases, all light entering the aperture is collected. The second case models the collection of emission from a molecule that is excited repeatedly; after each excitation, it rotates to a random orientation before emitting. These results are applicable to single-molecule detection techniques employing microdroplets and to other techniques for characterizing microparticles through the use of luminescence or inelastic scattering.

Contents

1	Introduction	1
2	The Sphere Dipole Fields and Aperture	3
3	Collection of Emission from a Fixed-Orientation Dipole	6
4	Collection of Emission from a Rotating Dipole	11
5	Summary	14
	Acknowledgements	15
	Appendices	17
A	Computation of the \bar{P}_{nm} , $\bar{\pi}_{nm}$, and $\bar{\tau}_{nm}$ Functions	17
B	The Green Function	19
C	Evaluation of I^{dot} and I^{cross}	21
D	$\{P_D\}$ in a Homogeneous Medium	22
	References	25
	Distribution	31
	Report Documentation Page	33

1. Introduction

Collection of emission from molecules in, on, or near small particles is important for a variety of applications: e.g., detection of single fluorescent molecules in microspheres [1–6], characterization of particles or droplets using fluorescence [7,8] or Raman [9] emission (the particles may be inhomogeneous particles, as in a droplet containing viruses or bacteria), characterization of lasing or nonlinear emission processes [10] or energy transfer [11] in microspheres, determination of the orientation of molecules on the surface of a droplet [12,13], and measurements of cavity quantum electrodynamic effects in spheres [13–16]. Collection of light elastically scattered by inclusions inside droplets [17–19] is a related problem. The irradiance collected by a lens depends on the position of the molecule(s) or scatterer(s) within the particle, because light reflects and refracts at the particle surface.

Fluorescence and Raman emission from molecules inside spheres have been modeled assuming the molecule can be approximated as an oscillating dipole [20–23]. The fields of the dipole and the fields induced inside and outside the sphere are expanded in spherical wave functions, and the expansion coefficients are determined by enforcement of the boundary conditions at the sphere’s surface. Real molecules emit with a nonzero transition linewidth. One can approximate the emission over the linewidth by averaging the output over a lineshape function [22].

In earlier work [22,23], Hill et al calculated the collection of fluorescence from a molecule, modeled as an oscillating dipole, using numerical integration over a circular collection aperture. In experimental work, it is often important to maximize the collection solid angle in order to maximize the signal. A problem arises when one uses numerical integration over the aperture: as the solid angle of the collection aperture increases, the number of points needed for accurate numerical integration can become prohibitive; the problem becomes worse as the size of the sphere increases. Problems with integrating over large-solid-angle apertures become even more important in modeling fluorescence or other emission in which the intensity must be integrated over emission frequency [22,23], particularly when high Q morphology-dependent resonances of the spheres are within the emission band.

Methods have been developed for “analytically” integrating, over a circular aperture, the scattering from a sphere illuminated with one or more plane waves [24–30]. In those methods, the integration computations are performed with recursion relations replacing the numerical integration scheme. Since numerical integration is not required, the approach is referred to as “analytical.” Advantages of the analytical method are that numerical results do not have to be checked for convergence (as they must be when numerical integration is used), and the analytical method can readily treat solid angles up to 4π sr.

In this report, we show that the emission from a dipole inside a spherical particle can be integrated analytically over a collection aperture. We consider two cases: (1) emission from a dipole with fixed orientation, and (2) emission from a dipole that rotates through all orientations. For case (2), we show how the irradiance collected by the aperture can be obtained by analytical integration over all dipole orientations. The rotating dipole model is appropriate for collection of fluorescent emission from a small molecule (e.g., laser dye) in water, where the molecule is excited repeatedly. In integrating the emitted irradiance over the circular aperture, we apply some of the techniques used previously for scattering by spheres [24–30].

In section 2, we describe the situation to be modeled: the sphere, dipole, electric fields, aperture, and coordinate systems. In section 3, we integrate the irradiance over the circular aperture when the dipole is in a fixed orientation, and provide (in eq (31)) a key result of this report: an explicit expression for the power collected. In section 4, we integrate the collected intensity at the aperture for a rotating dipole, and provide the results in equation (43). We summarize our work in section 5. Four appendices provide mathematical details.

2. The Sphere Dipole Fields and Aperture

In this section we define the problem (sphere, dipole, aperture, fields) for which we derive analytic solutions in sections 3 and 4. The region inside the sphere, region 1 with refractive index N_1 , is assumed to be homogeneous, except for one or more oscillating dipoles. The effect of any dipole (molecule) on the absorption of emission from other dipoles is included in the homogeneous refractive index of the sphere. The region outside the sphere, region 2 with refractive index N_2 , is infinite and homogeneous, except for the circular aperture and lens, which are far from the sphere. The wavenumbers are $k_0 = \omega/c$ in vacuum, $k_1 = N_1 k_0$ in region 1, and $k_2 = N_2 k_0$ in region 2.

An (x, y, z) Cartesian coordinate system with basis vectors $(\hat{\mathbf{x}}, \hat{\mathbf{y}}, \hat{\mathbf{z}})$ is located with its origin at the center of the sphere. The spherical system associated with the (x, y, z) Cartesian system is the (r, θ, ϕ) system with basis vectors $(\hat{\mathbf{r}}, \hat{\boldsymbol{\theta}}, \hat{\boldsymbol{\phi}})$. The z -axis, by definition, extends through the center of the circular aperture. An oscillating dipole \mathbf{p} , with $\exp(-i\omega t)$ time variation, is located at \mathbf{r}_d , and the orientation of the x -axis can be defined with respect to \mathbf{r}_d . We do this by giving the results in a form that holds for any chosen orientation of the x -axis.

The coordinates of \mathbf{r}_d in the spherical system are (r_d, θ_d, ϕ_d) . The half-plane containing the z -axis and \mathbf{r}_d is at an angle of ϕ_d from the x -axis, and the choice of ϕ_d has the effect of locating the x -axis with respect to \mathbf{r}_d . We give the results in a general form that holds for any value of ϕ_d .

Typically, an incident electromagnetic wave would be required to excite the dipole [20] (molecule), although there are exceptions: e.g., chemiluminescence and emission of thermal photons. We do not treat the problem of excitation of the dipole here. Methods of calculating the internal fields of spheres illuminated with plane waves [20,31,32], counterpropagating plane waves [23], or Gaussian beams [33,34] are well understood. For a molecule with a rotational diffusion time that is short compared to its average emission time, the excitation and emission problems can be separated (as described elsewhere [23]).

The electric field produced at \mathbf{r} in region 2 by dipole \mathbf{p} at \mathbf{r}_d in region 1 can be expressed as

$$\mathbf{E}(\mathbf{r}) = \omega^2 \mu_2 \mathbf{G}(\mathbf{r}, \mathbf{r}_d) \cdot \mathbf{p}(\mathbf{r}_d), \quad (1)$$

where $\mathbf{G}(\mathbf{r}, \mathbf{r}_d)$ is a dyadic Green function [35] for a source inside a sphere,* written here as

$$\begin{aligned} \mathbf{G}(\mathbf{r}, \mathbf{r}_d) = & \frac{ik_2\mu_2}{4\pi\mu_1} \sum_{n=1}^{\infty} \frac{2}{n(n+1)} \sum_{m=-n}^n (-1)^m \left[c_n \overline{\mathbf{M}}_{nm}^{(3)}(k_2\mathbf{r}) \overline{\mathbf{M}}_{n,-m}^{(1)}(k_1\mathbf{r}_d) \right. \\ & \left. + d_n \overline{\mathbf{N}}_{nm}^{(3)}(k_2\mathbf{r}) \overline{\mathbf{N}}_{n,-m}^{(1)}(k_1\mathbf{r}_d) \right]. \end{aligned} \quad (2)$$

The c_n and d_n are the internal field coefficients for region 1 as defined by Bohren and Huffman [31]. The $\overline{\mathbf{M}}_{nm}^{(q)}$ and $\overline{\mathbf{N}}_{nm}^{(q)}$ are the complex normalized vector spherical harmonics [36]

$$\overline{\mathbf{M}}_{nm}^{(q)}(k\mathbf{r}) \equiv \nabla \times \left(\mathbf{r} z_n^{(q)}(\rho) \exp(im\phi) \overline{P}_{nm}(\cos\theta) \right) \quad (3a)$$

$$= iz_n^{(q)}(\rho) \exp(im\phi) \left[\hat{\boldsymbol{\theta}} \pi_{nm}(\cos\theta) + i\hat{\boldsymbol{\phi}} \tau_{nm}(\cos\theta) \right], \quad (3b)$$

$$\overline{\mathbf{N}}_{nm}^{(q)}(k\mathbf{r}) \equiv \frac{1}{k} \nabla \times \overline{\mathbf{M}}_{nm}^{(q)}(k\mathbf{r}) \quad (4a)$$

$$\begin{aligned} = & \exp(im\phi) \left[\hat{\mathbf{r}} n(n+1) \frac{z_n^{(q)}(\rho)}{\rho} \overline{P}_{nm}(\cos\theta) \right. \\ & \left. + \frac{(d/d\rho) \left(\rho z_n^{(q)}(\rho) \right)}{\rho} \left(\hat{\boldsymbol{\theta}} \tau_{nm}(\cos\theta) + i\hat{\boldsymbol{\phi}} \pi_{nm}(\cos\theta) \right) \right], \end{aligned} \quad (4b)$$

where $\rho \equiv kr$, the $z_n^{(q)}(kr)$ are spherical Bessel functions $j_n(kr)$ when $q = 1$, and are spherical Hankel functions $h_n^{(1)}(kr)$ when $q = 3$, and where

$$\pi_{nm}(\cos\theta) = \frac{m \overline{P}_{nm}(\cos\theta)}{\sin\theta}, \quad (5)$$

$$\tau_{nm}(\cos\theta) = (d/d\theta) \overline{P}_{nm}(\cos\theta), \quad (6)$$

where the $\overline{P}_{nm}(\cos\theta)$ are the normalized associated Legendre functions of Arfken [37] and Belousov [38]. Computation of these functions is discussed in appendix A.

*The Green function shown here can be obtained following the approach of Tai [35]. The notation of equation (1) differs from Tai's notation in that it employs the complex vector spherical harmonics (see app B). Tai does the problem for a dipole outside a sphere. We used his approach to do the problem for a dipole inside a sphere.

The dipole is $\mathbf{p} = p\hat{\mathbf{p}}$, where p is the dipole magnitude and $\hat{\mathbf{p}}$ is a unit vector in the direction of the dipole axis. The dipole direction $\hat{\mathbf{p}}$ is specified by the Euler angles α and β (the ϕ and θ angles describing the dipole direction with respect to the (x, y, z) system), such that

$$\hat{\mathbf{p}} = \sum_{j=1}^3 f_j \hat{\mathbf{e}}_j, \quad (7)$$

where $(\hat{\mathbf{e}}_1, \hat{\mathbf{e}}_2, \hat{\mathbf{e}}_3) = (\hat{\mathbf{x}}, \hat{\mathbf{y}}, \hat{\mathbf{z}})$ and

$$f_1 = \sin \beta \cos \alpha, \quad (8a)$$

$$f_2 = \sin \beta \sin \alpha, \quad (8b)$$

$$f_3 = \cos \beta. \quad (8c)$$

Using equations (1), (2), and (7), we can write the electric field at \mathbf{r} generated by a dipole at \mathbf{r}_d as

$$\mathbf{E}(\mathbf{r}) = E_{ref} \sum_{n=1}^{\infty} \frac{2i}{n(n+1)} \sum_{m=-n}^n \sum_{j=1}^3 f_j \left[c_{n,-m}^j \overline{\mathbf{M}}_{nm}^{(3)}(k\mathbf{r}) + d_{n,-m}^j \overline{\mathbf{N}}_{nm}^{(3)}(k\mathbf{r}) \right], \quad (9)$$

where

$$E_{ref} \equiv \left(\frac{p}{4\pi\epsilon_0} \right) \left(\frac{(\mu_2)^2}{\mu_0\mu_1} \right) k_2(k_0)^2, \quad (10)$$

$$c_{n,-m}^j \equiv (-1)^m c_n \hat{\mathbf{e}}_j \cdot \overline{\mathbf{M}}_{n,-m}^{(1)}(k_1\mathbf{r}_d), \quad (11)$$

$$d_{n,-m}^j \equiv (-1)^m d_n \hat{\mathbf{e}}_j \cdot \overline{\mathbf{N}}_{n,-m}^{(1)}(k_1\mathbf{r}_d). \quad (12)$$

All the light emitted into the solid angle intercepted by the aperture is assumed to be collected. The aperture extends from θ_{\min} to θ_{\max} and is circular ($0 \leq \phi \leq 2\pi$). The solid angle with $\theta \leq \theta_{\min}$ is covered by a concentric aperture stop. With $d\Omega = \sin \theta \, d\theta \, d\phi$, the aperture solid angle is

$$\Delta\Omega = \int_0^{2\pi} \int_{\theta=\theta_{\min}}^{\theta=\theta_{\max}} d\Omega = 2\pi [\cos(\theta_{\min}) - \cos(\theta_{\max})]. \quad (13)$$

3. Collection of Emission from a Fixed-Orientation Dipole

The radiant power collected by the aperture is

$$P = \int_{\Delta\Omega} d\Omega r^2 I(\mathbf{r}), \quad (14)$$

where the irradiance of the emitted light at \mathbf{r} is

$$I(\mathbf{r}) = \frac{N_2}{2\mu_2 c} |\mathbf{E}(\mathbf{r})|^2. \quad (15)$$

Here, we use a dimensionless power collected by the aperture, P_D , defined as

$$P_D \equiv \frac{(k_2)^2}{I_{ref}} P, \quad (16)$$

where I_{ref} is the irradiance of a reference electromagnetic wave,

$$I_{ref} = \frac{N_2}{2\mu_2 c} (E_{ref})^2, \quad (17a)$$

$$= \frac{1}{8\pi} \frac{1}{4\pi\epsilon_0} \frac{(\mu_2)^3}{\mu_0(\mu_1)^2} c p^2 (k_2)^3 (k_0)^3. \quad (17b)$$

In terms of the electric field, the dimensionless power collected is

$$P_D = \int_{\Delta\Omega} d\Omega (k_2)^2 r^2 \frac{|\mathbf{E}(\mathbf{r})|^2}{(E_{ref})^2}. \quad (18)$$

The aperture is in the far field of the sphere. As $\rho_2 \equiv k_2 r$ becomes large, $h_n^{(1)}(\rho_2)$ can be replaced by its asymptotic expression, $(-i)^n [\exp(i\rho_2)]/i\rho_2$, and the $\overline{\mathbf{M}}_{nm}^{(3)}(k_2 \mathbf{r})$ and $\overline{\mathbf{N}}_{nm}^{(3)}(k_2 \mathbf{r})$ from equations (3b) and (4b) can be written

$$\begin{pmatrix} \overline{\mathbf{M}}_{nm}^{(3)}(k_2 \mathbf{r}) \\ \overline{\mathbf{N}}_{nm}^{(3)}(k_2 \mathbf{r}) \end{pmatrix} = (-i)^n \frac{\exp(i\rho_2)}{\rho_2} \exp(im\phi) \left[\hat{\theta} \begin{pmatrix} \overline{\pi}_{nm}(\cos \theta) \\ \overline{\tau}_{nm}(\cos \theta) \end{pmatrix} + i\hat{\phi} \begin{pmatrix} \overline{\tau}_{nm}(\cos \theta) \\ \overline{\pi}_{nm}(\cos \theta) \end{pmatrix} \right]. \quad (19)$$

Because $I(\mathbf{r})$ is inversely proportional to $(\rho_2)^2$ in the far field, P_D is not a function of r , the distance from the sphere's center to a field point \mathbf{r} on the aperture.

Substituting equation (9) for $\mathbf{E}(\mathbf{r})$ into equation (18), simplifying the result with the help of equation (19), and then integrating over ϕ from 0 to 2π , leads to

$$P_D = 8\pi \sum_{n=1}^{\infty} \sum_{n'=1}^{\infty} \sum_{m=-\min(n,n')}^{\min(n,n')} \sum_{j=1}^3 \sum_{j'=1}^3 f_j f_{j'} \left[\left(\psi_{nn',-m}^{dot} \right)^{jj'} \bar{\mathbf{I}}_{nn'm}^{dot} + \left(\psi_{nn',-m}^{cross} \right)^{jj'} \bar{\mathbf{I}}_{nn'm}^{cross} \right], \quad (20)$$

where

$$\left(\psi_{nn',-m}^{dot} \right)^{jj'} \equiv \frac{\left(c_{n,-m}^j c_{n',-m}^{j'*} + d_{n,-m}^j d_{n',-m}^{j'*} \right) (-i)^n (i)^{n'}}{n(n+1)n'(n'+1)}, \quad (21)$$

$$\left(\psi_{nn',-m}^{cross} \right)^{jj'} \equiv \frac{\left(c_{n,-m}^j d_{n',-m}^{j'*} + d_{n,-m}^j c_{n',-m}^{j'*} \right) (-i)^n (i)^{n'}}{n(n+1)n'(n'+1)}, \quad (22)$$

and

$$\begin{aligned} \bar{\mathbf{I}}_{nn'm}^{cross} &\equiv \int_{\mu_{\max}}^{\mu_{\min}} d\mu \left(\bar{\pi}_{nm}(\mu) \bar{\tau}_{n'm}(\mu) + \bar{\tau}_{nm}(\mu) \bar{\pi}_{n'm}(\mu) \right), \\ &= \left[-m \bar{P}_{nm}(\mu) \bar{P}_{n'm}(\mu) \right]_{\mu_{\max}}^{\mu_{\min}}, \end{aligned} \quad (23)$$

where $\mu \equiv \cos \theta$, $\mu_{\min} \equiv \cos \theta_{\min}$, and $\mu_{\max} \equiv \cos \theta_{\max}$, and

$$\bar{\mathbf{I}}_{nn'm}^{dot} \equiv \int_{\mu_{\max}}^{\mu_{\min}} d\mu \left(\bar{\pi}_{nm}(\mu) \bar{\pi}_{n'm}(\mu) + \bar{\tau}_{nm}(\mu) \bar{\tau}_{n'm}(\mu) \right). \quad (24)$$

The evaluation of $\bar{\mathbf{I}}_{nn'm}^{dot}$ requires consideration of two cases. When $n \neq n'$,

$$\bar{\mathbf{I}}_{nn'm}^{dot} = \left[(-1) \frac{[n(n+1) \bar{P}_{nm}(\mu) \sqrt{1-\mu^2} \bar{\tau}_{n'm}(\mu) - n'(n'+1) \bar{P}_{n'm}(\mu) \sqrt{1-\mu^2} \bar{\tau}_{nm}(\mu)]}{n(n+1) - n'(n'+1)} \right]_{\mu_{\max}}^{\mu_{\min}}. \quad (25)$$

When $n = n'$, $\bar{\mathbf{I}}_{nnm}^{dot}$ is obtained with the relation

$$\bar{\mathbf{I}}_{nnm}^{dot} = \left[(-1) \bar{P}_{nm}(\mu) \sqrt{1-\mu^2} \bar{\tau}_{nm}(\mu) \right]_{\mu_{\max}}^{\mu_{\min}} + n(n+1) \bar{I}_{nnm}, \quad (26)$$

where

$$\begin{aligned}\bar{I}_{nnm} &\equiv \int_{\mu_{\max}}^{\mu_{\min}} d\mu \left(\bar{P}_{nm}(\mu) \right)^2, \\ &= \left[\frac{\bar{P}_{nm}(\mu) \sqrt{1 - \mu^2} \bar{P}_{n,m-1}(\mu)}{\sqrt{(n+m)(n-m+1)}} \right]_{\mu_{\max}}^{\mu_{\min}} + \bar{I}_{nn,m-1}.\end{aligned}\quad (27)$$

The evaluation of the above integrals is discussed in appendix C.

The expression for the power collected, equation (20), can be simplified further: First, we use the general result,

$$\sum_{m=-M}^M F_m = \sum_{m=0}^M \epsilon_m \frac{1}{2} (F_m + F_{-m}), \quad (28)$$

where ϵ_m is the Neumann function defined as $\epsilon_0 = 1$ and $\epsilon_m = 2$ for $m \neq 0$; second, we use the relations

$$\bar{\mathbf{I}}_{nn',-m}^{dot} = \bar{\mathbf{I}}_{nn'm}^{dot}, \quad (29a)$$

$$\bar{\mathbf{I}}_{nn',-m}^{cross} = -\bar{\mathbf{I}}_{nn'm}^{cross}. \quad (29b)$$

Equations (29) above are obtained with

$$\bar{P}_{n,-m}(\mu) \equiv (-1)^m \bar{P}_{nm}(\mu), \quad (30a)$$

$$\bar{\pi}_{n,-m}(\mu) = (-1)^{m+1} \bar{\pi}_{nm}(\mu), \quad (30b)$$

$$\bar{\tau}_{n,-m}(\mu) = (-1)^m \bar{\tau}_{nm}(\mu), \quad (30c)$$

which follow from the function definitions. Using equations (20) and (28) to (30), we obtain our final result for the normalized power analytically integrated over the aperture:

$$\begin{aligned}P_D &= 4\pi \sum_{n=1}^{\infty} \sum_{n'=1}^{\infty} \sum_{m=0}^{\min(n,n')} \epsilon_m \left[\bar{\mathbf{I}}_{nn'm}^{dot} \sum_{j=1}^3 \sum_{j'=1}^3 f_j f_{j'} \left[\left(\psi_{nn'm}^{dot} \right)^{jj'} + \left(\psi_{nn',-m}^{dot} \right)^{jj'} \right] \right. \\ &\quad \left. - \bar{\mathbf{I}}_{nn'm}^{cross} \sum_{j=1}^3 \sum_{j'=1}^3 f_j f_{j'} \left[\left(\psi_{nn'm}^{cross} \right)^{jj'} - \left(\psi_{nn',-m}^{cross} \right)^{jj'} \right] \right].\end{aligned}\quad (31)$$

The sums over $f_j f_{j'} (\psi_{nn'm}^{dot})^{jj'}$ and $f_j f_{j'} (\psi_{nn'm}^{cross})^{jj'}$ are

$$\begin{aligned} \sum_{j=1}^3 \sum_{j'=1}^3 f_j f_{j'} (\psi_{nn'm}^{dot})^{jj'} &= \frac{(-i)^n (i)^{n'}}{n(n+1)n'(n'+1)} \\ &\times \left[\left(\sum_j f_j c_{nm}^j \right) \left(\sum_{j'} f_{j'} c_{n'm}^{j'} \right)^* \right. \\ &\left. + \left(\sum_j f_j d_{nm}^j \right) \left(\sum_{j'} f_{j'} d_{n'm}^{j'} \right)^* \right], \end{aligned} \quad (32)$$

and

$$\begin{aligned} \sum_{j=1}^3 \sum_{j'=1}^3 f_j f_{j'} (\psi_{nn'm}^{cross})^{jj'} &= \frac{(-i)^n (i)^{n'}}{n(n+1)n'(n'+1)} \\ &\times \left[\left(\sum_j f_j c_{nm}^j \right) \left(\sum_{j'} f_{j'} d_{n'm}^{j'} \right)^* \right. \\ &\left. + \left(\sum_j f_j d_{nm}^j \right) \left(\sum_{j'} f_{j'} c_{n'm}^{j'} \right)^* \right], \end{aligned} \quad (33)$$

where the sums can be expanded as

$$\sum_{j=1}^3 f_j c_{nm}^j = (-1)^m c_n \left[\sum_{j=1}^3 f_j \hat{\mathbf{e}}_j \cdot \bar{\mathbf{M}}_{nm}^{(1)} \right] \quad (34a)$$

$$\begin{aligned} &= (-1)^m c_n j_n(\rho_d) i \exp(im\phi_d) \left[\left[\sin \beta \cos \theta_d \cos(\alpha - \phi_d) \right. \right. \\ &\quad \left. \left. - \cos \beta \sin \theta_d \right] \bar{\pi}_{nm}(\cos \theta_d) \right. \\ &\quad \left. + \sin \beta \sin(\alpha - \phi_d) i \bar{\tau}_{nm}(\cos \theta_d) \right], \end{aligned} \quad (34b)$$

and

$$\sum_{j=1}^3 f_j d_{nm}^j = (-1)^m d_n \left[\sum_{j=1}^3 f_j \hat{\mathbf{e}}_j \cdot \bar{\mathbf{N}}_{nm}^{(1)} \right] \quad (35a)$$

$$\begin{aligned} &= (-1)^m d_n \exp(im\phi_d) \left[\left[\sin \beta \cos \theta_d \cos(\alpha - \phi_d) - \cos \beta \sin \theta_d \right] \right. \\ &\quad \times n(n+1) \frac{j_n(\rho_d)}{\rho_d} P_{nm}(\cos \theta_d) + \left(\left[\sin \beta \cos \theta_d \cos(\alpha - \phi_d) \right. \right. \\ &\quad \left. \left. - \cos \beta \sin \theta_d \right] \bar{\tau}_{nm}(\cos \theta_d) \right) \\ &\quad \left. + \sin \beta \sin(\alpha - \phi_d) i \bar{\pi}_{nm}(\cos \theta_d) \times \left(\frac{1}{\rho_d} \frac{d(\rho_d j_n(\rho_d))}{d\rho_d} \right) \right]. \quad (35b) \end{aligned}$$

To evaluate the terms with $-m$, i.e., $(\psi_{nn',-m}^{dot})^{jj'}$ and $(\psi_{nn',-m}^{cross})^{jj'}$ in equation (31), we rewrite equations (32) to (35) with $m \rightarrow (-m)$. Then equations (30) are used to evaluate the rewritten versions of equations (34) and (35).

4. Collection of Emission from a Rotating Dipole

A fluorescent molecule in a low-viscosity liquid may, after absorbing a photon, rotate many times before emitting a photon. For example, the rotational diffusion time for R6G in water is a few picoseconds, while the average time before emission is ~ 3 ns. In cases where the collection time is much longer than 3 ns, many photons are emitted in all directions from a rapidly rotating molecule as the fluorescence is collected, and we can obtain the resulting signal by averaging over all orientations of the rotating molecule.

Any function $g(\alpha, \beta)$, where α and β are the Euler angles, can be averaged over orientations with the expression

$$\{g(\alpha, \beta)\} \equiv \frac{1}{4\pi} \int_{\alpha=0}^{2\pi} d\alpha \int_{\beta=0}^{2\pi} d\beta \sin \beta g(\alpha, \beta). \quad (36)$$

We substitute $f_j f_{j'}$ for $g(\alpha, \beta)$ in this equation, use the definitions of equations (8), and evaluate the integrals to give

$$\{f_j f_{j'}\} = \frac{1}{3} \delta_{jj'}, \quad (37)$$

where $\delta_{jj'}$ is the Kronecker delta function.

We obtain the portion of the dimensionless power (emitted by a randomly rotating dipole) that is collected by the aperture $\{P_D\}$, by averaging P_D (given in eq (31)) over all dipole orientations. By using equations (36) and (37), we find the result to be

$$\begin{aligned} \{P_D\} = & \frac{4\pi}{3} \sum_{n=1}^{\infty} \sum_{n'=1}^{\infty} \sum_{m=0}^{\min(n,n')} \epsilon_m \left[\bar{\mathbf{I}}_{nn'm}^{dot} \sum_{j=1}^3 \left[\left(\psi_{nn'm}^{dot} \right)^{jj} + \left(\psi_{nn',-m}^{dot} \right)^{jj} \right] \right. \\ & \left. - \bar{\mathbf{I}}_{nn'm}^{cross} \sum_{j=1}^3 \left[\left(\psi_{nn'm}^{cross} \right)^{jj} - \left(\psi_{nn',-m}^{cross} \right)^{jj} \right] \right]. \end{aligned} \quad (38)$$

Evaluating the sums over j gives

$$\sum_{j=1}^3 \left(\psi_{nn'm}^{dot} \right)^{jj} = \sum_{j=1}^3 \left(\psi_{nn',-m}^{dot} \right)^{jj} = \Psi_{nn'm}^{dot}, \quad (39)$$

and

$$\sum_{j=1}^3 (\psi_{nn'm}^{cross})^{jj} = - \sum_{j=1}^3 (\psi_{nn',-m}^{cross})^{jj} = \Psi_{nn'm}^{cross} + \Psi_{n'nm}^{cross*}, \quad (40)$$

where

$$\begin{aligned} \Psi_{nn'm}^{dot} &\equiv \left(c_n c_{n'}^* \bar{\mathbf{M}}_{nm}^{(1)}(k\mathbf{r}_d) \cdot \bar{\mathbf{M}}_{n'm}^{(1)*}(k\mathbf{r}_d) + d_n d_{n'}^* \bar{\mathbf{N}}_{nm}^{(1)}(k\mathbf{r}_d) \cdot \bar{\mathbf{N}}_{n'm}^{(1)*}(k\mathbf{r}_d) \right) \\ &\quad \times \frac{(-i)^n (i)^{n'}}{n(n+1)n'(n'+1)}, \end{aligned} \quad (41)$$

$$\Psi_{nn'm}^{cross} \equiv \left(c_n d_{n'}^* \bar{\mathbf{M}}_{nm}^{(1)}(k\mathbf{r}_d) \cdot \bar{\mathbf{N}}_{n'm}^{(1)*}(k\mathbf{r}_d) \right) \frac{(-i)^n (i)^{n'}}{n(n+1)n'(n'+1)}. \quad (42)$$

The symmetry operation of interchanging the summation indices n and n' in $\Psi_{n'n,m}^{cross*}$ then allows equation (38) to be rewritten as

$$\{P_D\} = \frac{8\pi}{3} \sum_{n=1}^{\infty} \sum_{n'=1}^{\infty} \sum_{m=0}^{\min(n,n')} \epsilon_m \left[\Psi_{nn'm}^{dot} \bar{\mathbf{I}}_{nn'm}^{dot} - [2 \operatorname{Re}(\Psi_{nn'm}^{cross})] \bar{\mathbf{I}}_{nn'm}^{cross} \right]. \quad (43)$$

The radiant power of the collected emission from a sphere containing a rotating dipole is then computed with the relation

$$\{P\} = \frac{I_{ref}}{(k_2)^2} \{P_D\}. \quad (44)$$

In previous work [22,23], it was convenient to use the normalized fluorescence collected $F(\mathbf{r}_d, \omega)$, which is the fluorescence collected from a dipole in a sphere normalized by the fluorescence collected from a dipole in a homogeneous medium, i.e.,

$$F(\mathbf{r}_d, \omega) \equiv \frac{\{P\}}{\{P\}_{\text{hom}}}. \quad (45)$$

To calculate $\{P\}_{\text{hom}}$, we set the medium of the sphere identical to that of the surrounding medium (i.e., $N_1 = N_2$ and $\mu_1 = \mu_2$). We show in appendix D that $\{P_D\}_{\text{hom}} = (2/3)\Delta\Omega$, so that

$$\{P\}_{\text{hom}} = \left(\frac{I_{ref}}{(k_2)^2} \right) \left(\frac{2\Delta\Omega}{3} \right) \quad (46)$$

for any $\Delta\Omega$ and for any position of the dipole within the sphere. The normalized power collected by the aperture is then

$$F(\mathbf{r}_d, \omega) = \frac{4\pi}{\Delta\Omega} \sum_{n=1}^{\infty} \sum_{n'=1}^{\infty} \sum_{m=0}^{\min(n, n')} \epsilon_m \left[\Psi_{nn'm}^{dot} \bar{\mathbf{I}}_{nn'm}^{dot} - [2 \operatorname{Re} (\Psi_{nn'm}^{cross})] \bar{\mathbf{I}}_{nn'm}^{cross} \right]. \quad (47)$$

5. Summary

We have developed analytical expressions for the power entering a collecting aperture after emission from a dipole inside a spherical particle or droplet. The key results of this report are equations (31) and (43): equation (31) gives the power collected by an aperture when the dipole is in a fixed orientation, and equation (43) gives the power collected by the aperture when the emission is integrated over all dipole orientations.

The results should be useful in modeling experiments in which fluorescence or other emission from molecules inside, on, or near the surface of a sphere is measured. The results for integrating over dipole orientation are useful for modeling a dipole (at a fixed position) when the rotational diffusion time is much smaller than the fluorescence lifetime of the molecule, which in turn is much smaller than the collection time for the experiment; in such an experiment, the molecule absorbs and emits many photons from many different random orientations. An advantage of performing the integrations over the aperture analytically is that one need not test the results for convergence when the size, refractive index, or numerical aperture increase. We suggest that the results presented here may be particularly useful for cases in which the results must be integrated over a large wavelength range and the sphere is in a size range where narrow morphology-dependent resonances strongly influence the emission.

Acknowledgements

This research was sponsored in part by the U.S. Department of Energy, Office of Research and Development, under agreement DE-AI05-95OR22401 with the Army Research Laboratory. We thank Michael D. Barnes, William B. Whitten, and J. Michael Ramsey for insightful discussions and for stimulating our interest in the problems addressed here.

do not print

Appendix A. Computation of the \overline{P}_{nm} , $\overline{\pi}_{nm}$, and $\overline{\tau}_{nm}$ Functions

The $\overline{\pi}_{nm}(\mu)$ and $\overline{\tau}_{nm}(\mu)$ functions are defined here as

$$\overline{\pi}_{nm}(\mu) \equiv \frac{m\overline{P}_{nm}(\mu)}{\sqrt{1-\mu^2}}, \quad (\text{A-1})$$

$$\overline{\tau}_{nm}(\mu) \equiv -\sqrt{1-\mu^2}(d/d\mu)\overline{P}_{nm}(\mu), \quad (\text{A-2})$$

where $\overline{P}_{nm}(\mu)$ is the normalized associated Legendre function defined by Arfken [37] and Belousov [38] as

$$\overline{P}_{nm}(\mu) \equiv A_{nm}P_{nm}(\mu), \quad (\text{A-3})$$

where

$$A_{nm} \equiv \sqrt{\frac{(2n+1)(n-m)!}{2(n+m)!}}, \quad (\text{A-4})$$

and

$$P_{nm}(\mu) = (1-\mu^2)^{m/2} \frac{d^m}{d\mu^m} P_n(\mu) \quad (\text{A-5})$$

is the associated Legendre function as defined by Arfken [37] and Stratton [39] and used in the standard light scattering references by Bohren and Huffman [31] and van de Hulst [40]. These Legendre functions differ by a factor of $(-1)^m$ from the Legendre functions used by Jackson [41], Abramowitz and Stegun [42], and Gradshteyn and Ryzhik [43]. These last two references are of particular interest for their extensive recursion relations, but care must be exercised because of the $(-1)^m$ factor difference. The $\overline{\pi}_{nm}$ and $\overline{\tau}_{nm}$ functions are defined here to agree with Fuller's definitions [44], except with an additional factor of A_{nm} . There is a computational advantage to using normalized Legendre functions, because the factorials in A_{nm} do not have to be computed separately but are incorporated into the recursion relations.

We can obtain two relations convenient for computing $\overline{P}_{nm}(\mu)$, $\overline{\pi}_{nm}(\mu)$, and $\overline{\tau}_{nm}(\mu)$ by rewriting equations (8.733) given by Gradshteyn and Ryzhik [43] as

$$\begin{aligned} \mu \overline{\pi}_{nm}(\mu) = & \left(\frac{1}{2} \right) \left[\sqrt{(n-m+1)(n+m)} \overline{P}_{n,m-1}(\mu) \right. \\ & \left. + \sqrt{(n+m+1)(n-m)} \overline{P}_{n,m+1}(\mu) \right], \end{aligned} \quad (\text{A-6})$$

$$\begin{aligned} \overline{\tau}_{nm}(\mu) = & \left(\frac{1}{2} \right) \left[\sqrt{(n-m+1)(n+m)} \overline{P}_{n,m-1}(\mu) \right. \\ & \left. - \sqrt{(n+m+1)(n-m)} \overline{P}_{n,m+1}(\mu) \right]. \end{aligned} \quad (\text{A-7})$$

If $\mu \neq 1$, then equation (A-1) is used to compute $\overline{\pi}_{nm}(\mu)$ from $\overline{P}_{nm}(\mu)$, and the above two relations are used to compute $\overline{P}_{n,m+1}(\mu)$ and $\overline{\tau}_{nm}(\mu)$.

If $\mu = 1$, then

$$\overline{P}_{nm}(1) = \sqrt{\frac{(2n+1)}{2}} \delta_{m,0}, \quad (\text{A-8})$$

and equations (A-6) and (A-7) become

$$\overline{\pi}_{nm}(1) = \frac{1}{2} \sqrt{\frac{(2n+1)n(n+1)}{2}} (\delta_{m,1} + \delta_{m,-1}), \quad (\text{A-9})$$

$$\overline{\tau}_{nm}(1) = \frac{1}{2} \sqrt{\frac{(2n+1)n(n+1)}{2}} (\delta_{m,1} - \delta_{m,-1}). \quad (\text{A-10})$$

Appendix B. The Green Function

Using the traditional vector spherical harmonics, Tai [35] writes the equations necessary to solve for the Green functions when the dipole is outside the sphere. Tai [35] does not explicitly give the results for his coefficients A_n , B_n , C_n , and D_n . However, when his equations are solved, his upper case coefficients can be seen to be related to the lower case sphere coefficients of Bohren and Huffman [31], as

$$\begin{aligned} A_n &= -b_n, \\ B_n &= -a_n, \\ C_n &= c_n, \\ D_n &= d_n. \end{aligned} \tag{B-1}$$

The Green functions for a dipole inside the sphere can be obtained with essentially the same approach used by Tai [35] for the external dipole, and we obtain the result for the region outside the sphere as

$$\begin{aligned} \mathbf{G}(\mathbf{r}, \mathbf{r}_d) &= \frac{ik_2\mu_2}{4\pi\mu_1} \sum_{n=1}^{\infty} \frac{2}{n(n+1)} \sum_{m=0}^n \epsilon_m (A_{nm})^2 \sum_{\sigma=e,o} \left[c_n \mathbf{M}_{\sigma mn}^{(3)}(k_2\mathbf{r}) \mathbf{M}_{\sigma mn}^{(1)}(k_1\mathbf{r}_d) \right. \\ &\quad \left. + d_n \mathbf{N}_{\sigma mn}^{(3)}(k_2\mathbf{r}) \mathbf{N}_{\sigma mn}^{(1)}(k_1\mathbf{r}_d) \right]. \end{aligned} \tag{B-2}$$

To rewrite this result with our vector spherical harmonics (eq (3b) and (4b)), we note that they are related to the traditional ones [31,32,39] by the relations

$$A_{nm} \mathbf{M}_{emn}^{(q)} = \frac{1}{2} \left(\overline{\mathbf{M}}_{nm}^{(q)} + (-1)^m \overline{\mathbf{M}}_{n,-m}^{(q)} \right), \tag{B-3}$$

$$A_{nm} \mathbf{M}_{omn}^{(q)} = \frac{1}{2i} \left(\overline{\mathbf{M}}_{nm}^{(q)} - (-1)^m \overline{\mathbf{M}}_{n,-m}^{(q)} \right), \tag{B-4}$$

$$A_{nm}^{(q)} \mathbf{N}_{emn} = \frac{1}{2} \left(\overline{\mathbf{N}}_{nm}^{(q)} + (-1)^m \overline{\mathbf{N}}_{n,-m}^{(q)} \right), \tag{B-5}$$

$$A_{nm} \mathbf{N}_{omn}^{(q)} = \frac{1}{2i} \left(\overline{\mathbf{N}}_{nm}^{(q)} - (-1)^m \overline{\mathbf{N}}_{n,-m}^{(q)} \right), \tag{B-6}$$

where A_{nm} is defined in equation (A-4).

Using equations (B-3) to (B-6) we rewrite the Green function as

$$\begin{aligned}
\mathbf{G}(\mathbf{r}, \mathbf{r}_d) = & \frac{ik_2\mu_2}{4\pi\mu_1} \sum_{n=1}^{\infty} \frac{1}{n(n+1)} \sum_{m=0}^n \epsilon_m (-1)^m \\
& \times \left[c_n \left(\overline{\mathbf{M}}_{nm}^{(3)}(k_2\mathbf{r}) \overline{\mathbf{M}}_{n,-m}^{(1)}(k_1\mathbf{r}_d) + \overline{\mathbf{M}}_{n,-m}^{(3)}(k_2\mathbf{r}) \overline{\mathbf{M}}_{nm}^{(1)}(k_1\mathbf{r}_d) \right) \right. \\
& \left. + d_n \left(\overline{\mathbf{N}}_{nm}^{(3)}(k_2\mathbf{r}) \overline{\mathbf{N}}_{n,-m}^{(1)}(k_1\mathbf{r}_d) + \overline{\mathbf{N}}_{n,-m}^{(3)}(k_2\mathbf{r}) \overline{\mathbf{N}}_{nm}^{(1)}(k_1\mathbf{r}_d) \right) \right]. \quad (\text{B-7})
\end{aligned}$$

Then using the general relation for sums given in equation (28), we obtain

$$\begin{aligned}
\mathbf{G}(\mathbf{r}, \mathbf{r}_d) = & \frac{ik_2\mu_2}{4\pi\mu_1} \sum_{n=1}^{\infty} \frac{2}{n(n+1)} \sum_{m=-n}^n (-1)^m \\
& \times \left[c_n \overline{\mathbf{M}}_{nm}^{(3)}(k_2\mathbf{r}) \overline{\mathbf{M}}_{n,-m}^{(1)}(k_1\mathbf{r}_d) + d_n \overline{\mathbf{N}}_{nm}^{(3)}(k_2\mathbf{r}) \overline{\mathbf{N}}_{n,-m}^{(1)}(k_1\mathbf{r}_d) \right]. \quad (\text{B-8})
\end{aligned}$$

Appendix C. Evaluation of I^{dot} and I^{cross}

The result for $\bar{\mathbf{I}}_{nn'm}^{cross}$ given in equation (23) in the body of the report is obtained by inspection.

We can prove the expressions for $\bar{\mathbf{I}}_{nn'm}^{dot}$ (eq (25) to (27)) by taking the derivative with respect to μ of both sides of the equations, and then using equations (A-1) and (A-2). Equation (27) can be proved with the help of

$$\bar{P}_{n,m-1}(\mu) = \frac{\mu \bar{\pi}_{nm}(\mu) + \bar{\tau}_{nm}(\mu)}{\sqrt{(n+m)(n-m+1)}}, \quad (\text{C-1})$$

which we obtain by adding equations (A-6) and (A-7). It is also helpful to know that the differential equation for the associated Legendre function can be written as

$$\frac{d}{d\mu} \left(\sqrt{1-\mu^2} \bar{\tau}_{nm}(\mu) \right) = n(n+1) \bar{P}_{nm}(\mu) - \frac{m \bar{\pi}_{nm}(\mu)}{\sqrt{1-\mu^2}}. \quad (\text{C-2})$$

Appendix D. $\{P_D\}$ in a Homogeneous Medium

To check the derivations and to obtain a value for normalization, we set the medium of the sphere identical to the medium surrounding the sphere (i.e., $N_1 = N_2$ and $\mu_1 = \mu_2$, where both N_1 and N_2 are real). Then equation (43) can be shown to give the result $\{P_D\} = (2/3)\Delta\Omega$ for any $\Delta\Omega$ and for any position of the dipole within the sphere. We start with a proof for $\Delta\Omega = 4\pi$, and extend that result with an intuitive argument to arbitrary $\Delta\Omega$.

If $N_1 = N_2$ and $\mu_1 = \mu_2$, then $c_n = d_n = 1$. Further, if $\Delta\Omega = 4\pi$, then $\bar{\mathbf{I}}_{nn'm}^{dot} = n(n+1)\delta_{nn'}$ and $\bar{\mathbf{I}}_{nn'm}^{cross} = 0$, so that equation (43) becomes

$$\{P_D\}_{\text{hom}}^{\Delta\Omega=4\pi} = \frac{8\pi}{3} \sum_{n=1}^{\infty} \sum_{m=0}^n \frac{\epsilon_m (|\bar{\mathbf{M}}_{nm}^{(1)}(k_1 \mathbf{r}_d)|^2 + |\bar{\mathbf{N}}_{nm}^{(1)}(k_1 \mathbf{r}_d)|^2)}{n(n+1)}. \quad (\text{D-1})$$

Because N_1 is real here, $\rho_d \equiv k_1 r_d$ is also real. This allows the $|\bar{\mathbf{M}}_{nm}^{(1)}(k_1 \mathbf{r}_d)|^2$ and $|\bar{\mathbf{N}}_{nm}^{(1)}(k_1 \mathbf{r}_d)|^2$ to be written as

$$|\bar{\mathbf{M}}_{nm}^{(1)}(k_1 \mathbf{r}_d)|^2 = (j_n(\rho_d))^2 \left([\bar{\pi}_{nm}(\cos \theta_d)]^2 + [\bar{\tau}_{nm}(\cos \theta_d)]^2 \right), \quad (\text{D-2})$$

$$\begin{aligned} |\bar{\mathbf{N}}_{nm}^{(1)}(k_1 \mathbf{r}_d)|^2 &= n^2(n+1)^2 \left(\frac{j_n(\rho_d)}{\rho_d} \right)^2 \left(\bar{P}_{nm}(\cos \theta_d) \right)^2 \\ &\quad + \left(\frac{(d/d\rho_d)(\rho_d j_n(\rho_d))}{\rho_d} \right)^2 \left([\bar{\pi}_{nm}(\cos \theta_d)]^2 + [\bar{\tau}_{nm}(\cos \theta_d)]^2 \right). \end{aligned} \quad (\text{D-3})$$

The addition theorem for spherical harmonics given by Arfken (eq (12.197), p 694) [37] can be rewritten as

$$\sum_{m=0}^n \epsilon_m \left(\bar{P}_{nm}(\cos \theta_d) \right)^2 = \frac{2n+1}{2}, \quad (\text{D-4})$$

and a summation over vector-spherical harmonics given by Jackson (eq (16.77), p 753) [41] can be rewritten as

$$\sum_{m=0}^n \epsilon_m \left[(\bar{\pi}_{nm}(\cos \theta_d))^2 + (\bar{\tau}_{nm}(\cos \theta_d))^2 \right] = \frac{(2n+1)n(n+1)}{2}, \quad (\text{D-5})$$

so that

$$\begin{aligned} \{P_D\}_{\text{hom}}^{\Delta\Omega=4\pi} = & \frac{4\pi}{3} \sum_{n=1}^{\infty} (2n+1) \left[\left(j_n(\rho_d) \right)^2 + n(n+1) \left(\frac{j_n(\rho_d)}{\rho_d} \right)^2 \right. \\ & \left. + \left(\frac{(d/d\rho_d)(\rho_d j_n(\rho_d))}{\rho_d} \right)^2 \right]. \end{aligned} \quad (\text{D-6})$$

The Rayleigh equation (Arfken [37], p 665) gives

$$\exp(i\rho_d\mu) = \sum_{n=0}^{\infty} i^n (2n+1) j_n(\rho_d) P_n(\mu), \quad (\text{D-7})$$

where $P_n(\mu)$ is a Legendre polynomial. Multiplying each side of equation (D-7) by its complex conjugate and integrating over μ from -1 to 1 gives

$$\sum_{n=1}^{\infty} (2n+1) (j_n(\rho_d))^2 = 1 - (j_0(\rho_d))^2. \quad (\text{D-8})$$

The Rayleigh equation can be used in essentially the same way to obtain the result

$$\sum_{n=1}^{\infty} (2n+1) \left(\frac{(d/d\rho_d)(\rho_d j_n(\rho_d))}{\rho_d} \right)^2 = \frac{1}{3} + \frac{1}{(\rho_d)^2} - \left(\frac{(d/d\rho_d)(\rho_d j_0(\rho_d))}{\rho_d} \right)^2. \quad (\text{D-9})$$

By using the differential equation of the Legendre polynomial and the Rayleigh equation (D-7), and then multiplying and integrating as before, one can obtain the result

$$\sum_{n=1}^{\infty} (2n+1) n(n+1) \left(\frac{j_n(\rho_d)}{\rho_d} \right)^2 = \frac{2}{3}. \quad (\text{D-10})$$

Recalling that $j_0(\rho_d) = [\sin(\rho_d)]/\rho_d$ then allows these three sums to be combined to give the exact result

$$\{P_D\}_{\text{hom}}^{\Delta\Omega=4\pi} = \frac{8\pi}{3}. \quad (\text{D-11})$$

Because the emission from a randomly rotating dipole in a homogeneous medium is isotropic in the far field, the power $\{P_D\}$ collected by an aperture solid angle $\Delta\Omega$ in the far field is simply proportional to $\Delta\Omega$. Then

$$\begin{aligned} \{P_D\}_{\text{hom}} &= \left(\frac{8\pi}{3} \right) \left(\frac{\Delta\Omega}{4\pi} \right) \\ &= \frac{2}{3} \Delta\Omega \end{aligned} \quad (\text{D-12})$$

is an exact result for all $\Delta\Omega$.

do not print

References

1. M. D. Barnes, W. B. Whitten, and J. M. Ramsey, "Detecting single molecules in liquids," *Anal. Chem.* **67**, 418A–423A (1995).
2. M. D. Barnes, W. B. Whitten, and J. M. Ramsey, "Detection of single Rhodamine 6G molecules in levitated microdroplets," *Anal. Chem.* **65**, 2360–2365 (1993).
3. W. B. Whitten and J. M. Ramsey, "Single-molecule detection limits in levitated microdroplets," *Anal. Chem.* **63**, 1027–1031 (1991).
4. W. B. Whitten and J. M. Ramsey, "Photocount probability distributions for single fluorescent molecules," *Appl. Spectrosc.* **46**, 1587–1589 (1992).
5. K. C. Ng, W. B. Whitten, S. Arnold, and J. M. Ramsey, "Digital chemical analysis of dilute microdroplets," *Anal. Chem.* **64**, 2914–2919 (1992).
6. M. D. Barnes, C.-Y. Kung, W. B. Whitten, J. M. Ramsey, and S. Arnold, "Molecular fluorescence in a microcavity: Solvation dynamics and single molecule detection," *Optical Processes in Microcavities*, R. K. Chang and A. J. Campillo, eds. (World Scientific, Singapore, 1996), pp 135–165.
7. R. G. Pinnick, S. C. Hill, P. Nachman, J. D. Pendleton, G. L. Fernandez, M. W. Mayo, and J. G. Bruno, "Fluorescence particle counter for detecting airborne bacteria and other biological particles," *Aerosol Sci. Technol.* **23**, 653–664 (1995).
8. G. Chen, P. Nachman, R. G. Pinnick, S. C. Hill, and R. K. Chang, "Conditional-firing aerosol-fluorescence spectrum analyzer for individual airborne particles with pulsed 266-nm laser excitation," *Opt. Lett.* **21**, 1307–1309 (1996).
9. M. F. Buehler, T. M. Allen, and E. J. Davis, "Microparticle Raman spectroscopy of multicomponent aerosols," *J. Colloid Interface Sci.* **146**, 79 (1991).

10. J.-Z. Zhang, G. Chen, and R. K. Chang, "Pumping of stimulated Raman scattering by stimulated Brillouin scattering within a single liquid droplet: Input laser linewidth effects," *J. Opt. Soc. Am. B* **7**, 108–115 (1990). J. C. Swindal, D. H. Leach, R. K. Chang, and K. Young, "Precession of morphology-dependent resonances in non-spherical liquid droplets," *Opt. Lett.* **18**, 191–193 (1993). For other examples, see the review chapter by S. C. Hill and R. K. Chang, "Nonlinear optics in droplets," *Studies in Classical and Quantum Nonlinear Optics*, O. Keller, ed. (Nova Science, Commack, NY, 1995), pp 171–242.
11. S. D. Druger, S. Arnold, and L. M. Folan, "Theory of enhanced energy transfer between molecules embedded in spherical dielectric particles," *J. Chem. Phys.* **87**, 2649 (1987).
12. L. Folan and S. Arnold, "Determination of molecular orientation at the surface of an aerosol particle by morphology-dependent photoselection," *Opt. Lett.* **13**, 1–3 (1988).
13. M. D. Barnes, C.-Y. Kung, W. B. Whitten, J. M. Ramsey, S. Arnold, and S. Holler, "Fluorescence of oriented molecules in a microcavity," *Phys. Rev. Lett.* **76**, 3931–3934 (1996).
14. H.-B. Lin, J. D. Eversole, and A. J. Campillo, "Cavity-modified spontaneous emission rates in liquid microdroplets," *Phys. Rev. A* **45**, 6756 (1992).
15. M. D. Barnes, W. B. Whitten, S. Arnold, and J. M. Ramsey, "Homogeneous linewidths of Rhodamine 6G at room temperature from cavity-enhanced spontaneous emission rates," *J. Chem. Phys.* **97**, 7842–7845 (1992).
16. M. D. Barnes, W. B. Whitten, and J. M. Ramsey, "Enhanced fluorescence yields through cavity-QED effects in microdroplets," *J. Opt. Soc. Am. B* **11**, 1297–1304 (1994).
17. B. V. Bronk, M. J. Smith, and S. Arnold, "Photon-correlation spectroscopy for small spherical inclusions in a micrometer-sized electrody-namically levitated droplet," *Opt. Lett.* **18**, 93–95 (1993).
18. D. Ngo and R. G. Pinnick, "Suppression of scattering resonances in in-homogeneous microdroplets," *J. Opt. Soc. Am. A* **11**, 1352–1359 (1994).

19. S. C. Hill, H. I. Saleheen, and K. A. Fuller, "Volume current method for modeling light scattering by inhomogeneously perturbed spheres," *J. Opt. Soc. Am. A* **12**, 905–915 (1995).
20. H. Chew, P. J. McNulty, and M. Kerker, "Model for Raman and fluorescent scattering by molecules embedded in small particles," *Phys. Rev. A* **13**, 396–404 (1976).
21. S. Druger and P. J. McNulty, "Radiation patterns of fluorescence from molecules embedded in small particles: General case," *Appl. Opt.* **22**, 75–82 (1983).
22. S. C. Hill, H. I. Saleheen, M. D. Barnes, W. B. Whitten, and J. M. Ramsey, "Modeling fluorescence collection from single molecules in microspheres: Effects of position, orientation, and frequency," *Appl. Opt.* **35**, 6278–6288 (1996).
23. S. C. Hill, M. D. Barnes, W. B. Whitten, and J. M. Ramsey, "Collection of fluorescence from single molecules in microspheres: Effects of illumination geometry," *Appl. Opt.*, in press.
24. P. Chylek, "Mie scattering into the backward hemisphere," *J. Opt. Soc. Am.* **63**, 1467–1471 (1973).
25. W. J. Wiscombe and P. Chylek, "Mie scattering between any two angles," *J. Opt. Soc. Am.* **67**, 572–573 (1977).
26. W. P. Chu and D. M. Robinson, "Scattering from a moving spherical particle by two crossed coherent plane waves," *Appl. Opt.* **16**, 619–626 (1977).
27. J. D. Pendleton, "Mie scattering into apertures," *J. Opt. Soc. Am.* **72**, 1029–1033 (1982).
28. J. D. Pendleton, *A Generalized Mie Theory Solution and Its Application to Particle Sizing Interferometry*, Ph.D. thesis (University of Tennessee, Knoxville, 1982), p 90.
29. J. Y. Son, W. M. Farmer, and T. V. Giel, Jr., "New optical geometry for the particle sizing interferometer," *Appl. Opt.* **25**, 4332–4337 (1986).

30. J. Y. Son, *Multiple Methods for Obtaining Particle Size Distribution with a Particle Sizing Interferometer*, Ph.D. thesis (University of Tennessee, Knoxville, 1985).
31. C. F. Bohren and D. R. Huffman, *Absorption and Scattering of Light by Small Particles* (Wiley, New York, 1983), pp 82–101.
32. P. W. Barber and S. C. Hill, *Light Scattering by Particles: Computational Methods* (World Scientific, Singapore, 1990).
33. J. P. Barton and D. R. Alexander, “Electromagnetic fields for an irregularly shaped, near-spherical particle illuminated by a focused laser beam,” *J. Appl. Phys.* **69**, 7973–7986 (1991).
34. E.E.M. Khaled, S. C. Hill, and P. W. Barber, “Scattered and internal intensity of a sphere illuminated with a Gaussian beam,” *IEEE Trans. Antennas Propag.* **41**, 295–303 (1993).
35. C.-T. Tai, *Dyadic Green Functions in Electromagnetic Theory*, 2nd ed. (IEEE Press, Piscataway, NJ, 1994), ch. 10.
36. G. Videen, D. Ngo, P. Chylek, and R. G. Pinnick, “Light scattering from a sphere with an irregular inclusion,” *J. Opt. Soc. Am. A* **12**, 922–928 (1995).
37. G. Arfken, *Mathematical Methods for Physicists*, 3rd ed. (Academic Press, San Diego, 1985), pp 198–200, 253, 678.
38. S. L. Belousov, *Tables of Normalized Associated Legendre Polynomials* (Pergamon, New York, 1962).
39. J. A. Stratton, *Electromagnetic Theory* (McGraw-Hill, New York, 1941), p 401.
40. H. C. van de Hulst, *Light Scattering by Small Particles* (Dover, New York, 1981), pp 34, 124.
41. J. D. Jackson, *Classical Electrodynamics*, 2nd ed. (Wiley, New York, 1975), p 746.
42. M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions* (Dover, New York, 1964).

- 43. I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products* (Academic, New York, 1980), pp 1005, 1008.
- 44. K. A. Fuller, "Scattering and absorption cross sections of compounded spheres. I. Theory for external aggregation," *J. Opt. Soc. Am. A* **11**, 3251–3260 (1994).

Distribution

Admnstr
Defns Techl Info Ctr
Attn DTIC-OCF
8725 John J Kingman Rd Ste 0944
FT Belvoir VA 22060-6218

Central Intllgnc Agency Dir DB Standard
Attn GE 47 QB
Washington DC 20505

Chairman Joint Chiefs of Staff
Attn J5 R&D Div
Washington DC 20301

Defns Intllgnc Agency
Attn DT 2 Wpns & Sys Div
Washington DC 20301

Dir of Defns Rsrch & Engrg
Attn DD TWP
Attn Engrg
Washington DC 20301

Ofc of the Dir Rsrch and Engrg
Attn R Menz
Pentagon Rm 3E1089
Washington DC 20301-3080

Ofc of the Secy of Defns
Attn ODDRE (R&AT) G Singley
Attn ODDRE (R&AT) S Gontarek
The Pentagon
Washington DC 20301-3080

OIR CSB CRB
Attn A M Jones
RB 1413 OHM
Washington DC 20505

OSD
Attn OUSD(A&T)/ODDDR&E(R) J Lupo
The Pentagon
Washington DC 20301-7100

Commanding Officer
Attn NMCB23
6205 Stuart Rd Ste 101
FT Belvoir VA 22060-5275

CECOM
Attn PM GPS COL S Young
FT Monmouth NJ 07703

CECOM RDEC Elect System Div Dir
Attn J Niemela
FT Monmouth NJ 07703

CECOM
Sp & Terrestrial Commctn Div
Attn AMSEL-RD-ST-MC-M H Soicher
FT Monmouth NJ 07703-5203

Dir of Chem & Nuc Ops DA DCSOPS
Attn Techl Lib
Washington DC 20301

Dpty Assist Secy for Rsrch & Techl
Attn SARD-TT F Milton Rm 3E479
The Pentagon
Washington DC 20301-0103

Hdqtrs Dept of the Army
Attn DAMO-FDT D Schmidt Rm 3C514
400 Army Pentagon
Washington DC 20301-0460

US Army Engrg Div
Attn HNDED FD
PO Box 1500
Huntsville AL 35807

US Army ERDEC
Attn I Sindoni
Attn J Embury
Attn M Milham
Attn S Christesen
Attn S Godoff
Attn B Bronk
Aberdeen Proving Ground MD 21005-5423

US Army Matl Cmnd
Dpty CG for RDE Hdqtrs
Attn AMCRD BG Beauchamp
5001 Eisenhower Ave
Alexandria VA 22333-0001

US Army Matl Cmnd
Prin Dpty for Acquisition Hdqtrs
Attn AMCDCG-A D Adams
5001 Eisenhower Ave
Alexandria VA 22333-0001

Distribution (cont'd)

US Army Matl Cmnd
Prin Dpty for Techlgy Hdqrts
Attn AMCDCG-T M Fiset
5001 Eisenhower Ave
Alexandria VA 22333-0001

US Army Mis & Spc Intllgnc Ctr
Attn AIAMS YDL
Redstone Arsenal AL 35898-5500

US Army NGIC
Attn Rsrch & Data Branch
220 7th Stret NE
Charlottesville VA 22901-5396

US Army Nuc & Cheml Agency
7150 Heller Loop Ste 101
Springfield VA 22150-3198

US Army Rsrch Ofc
Attn AMXRO-ICA B Mann
PO Box 12211
Research Triangle Park NC 27709-2211

US Army Strtgc Defns Cmnd
Attn CSSD H MPL Techl Lib
Attn CSSD H XM Davies
PO Box 1500
Huntsville AL 35807

US Military Academy
Dept of Mathematical Sci
Attn MAJ D Engen
West Point NY 10996

USAASA
Attn MOAS-AI W Parron
9325 Gunston Rd Ste N319
FT Belvoir VA 22060-5582

Chief of Nav OPS Dept of the Navy
Attn OP 03EG
Washington DC 20350

GPS Joint Prog Ofc Dir
Attn COL J Clay
2435 Vela Way Ste 1613
Los Angeles AFB CA 90245-5500

Special Assist to the Wing Cmndr
Attn 50SW/CCX Capt P H Bernstein
300 O'Malley Ave Ste 20
Falcon AFB CO 80912-3020

DARPA
Attn B Kaspar
Attn L Stotts
Attn Techl Lib
3701 N Fairfax Dr
Arlington VA 22203-1714

US Dept of Energy
Attn KK 22 K Sisson
Attn Techl Lib
Washington DC 20585

ARL Electromag Group
Attn Campus Mail Code F0250 A Tucker
University of Texas
Austin TX 78712

US Army Rsrch Lab
Attn AMSRL-CI-LL Techl Lib (3 copies)
Attn AMSRL-CS-AL-TA Mail & Records
Mgmt
Attn AMSRL-CS-AL-TP Techl Pub (3 copies)
Attn AMSRL-IS-EE S C Hill (15 copies)
Attn AMSRL-IS-EE J D Pendleton (15 copies)
Adelphi MD 20783-1197

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE December 1997		3. REPORT TYPE AND DATES COVERED Final, from December 1995 to May 1997
4. TITLE AND SUBTITLE Collection of Emission from an Oscillating Dipole Inside a Sphere: Analytic Integration Over a Circular Aperture			5. FUNDING NUMBERS DA PR: B53A PE: P61102	
6. AUTHOR(S) J. David Pendleton and Steven C. Hill				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory Attn: AMSRL-IS-EE 2800 Powder Mill Road Adelphi, MD 20783-1197			8. PERFORMING ORGANIZATION REPORT NUMBER ARL-TR-1399	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory 2800 Powder Mill Road Adelphi, MD 20783-1197			10. SPONSORING/MONITORING AGENCY REPORT NUMBER U.S. DOE Office of Research and Development Washington, DC 20585	
11. SUPPLEMENTARY NOTES AMS code: 611102.53A11 ARL PR: 7FEJ60				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) We describe a method for analytically integrating, over a circular aperture, the emission from an oscillating dipole inside a dielectric sphere. The model is useful for investigating fluorescence, Raman, or other emission from molecules inside spherical particles or droplets. The analysis is performed for two cases: (1) the dipole emits from a fixed orientation, and (2) the dipole emits from all orientations. In both cases, all light entering the aperture is collected. The second case models the collection of emission from a molecule that is excited repeatedly; after each excitation, it rotates to a random orientation before emitting. These results are applicable to single-molecule detection techniques employing microdroplets and to other techniques for characterizing microparticles through the use of luminescence or inelastic scattering.				
14. SUBJECT TERMS Fluorescence, electromagnetic waves, emission, microspheres, scattering, single-molecule detection			15. NUMBER OF PAGES 35	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	